

Botox Consent

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances. However, informed consent documents should not be considered all-inclusive in defining other methods of care and risks encountered. Your physician may provide you with additional or different information which is based on all of the facts pertaining to your particular case and the current state of medical knowledge. Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

PATIENT INFORMATION FOR BOTOX® COSMETIC

Prior to BOTOX injection(s):

- If you are on a blood-thinner including aspirin or baby aspirin, this must be discussed with your physician one week prior to treatment. The risk of bruising and bleeding is significantly higher in patients that have blood-thinners in their system.

After BOTOX injection(s):

- Do not lie down for four hours after treatment.
- Do not massage the treated area. This can cause BOTOX to spread to the other muscles increasing the risks of complications.
- No strenuous exercise for 6 hours after treatment.
- BOTOX cosmetic maximum results are usually achieved about 10 days after treatment. Occasionally, a series of treatments are needed to receive the desired results. There will be a charge incurred for each treatment required to achieve those results. Injections can be repeated every 3-6 months.

INSTRUCTIONS

This is an informed-consent document which has been prepared to help your physician inform you concerning Botox (onabotulinumtoxinA) injection, its risks, and alternative treatments. It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for this procedure as proposed by your physician and agreed upon by you.

GENERAL INFORMATION

When Botox is injected, it produces a temporary paralysis of muscle by preventing transmission of nerve impulses to muscle. The duration of muscle paralysis generally lasts for approximately three to four months. Botox has been FDA approved to treat glabellar lines and crow's feet. Botox injections are customized for every patient, depending on his or her particular needs. Botox cannot stop the process of aging. It can however, temporarily diminish the look of wrinkles caused by muscle groups. Botox injections may be performed as a singular procedure or as an adjunct to a surgical procedure.

RISKS of Botox (Botulinum Toxin Type A) Injections

Every procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your physician to make sure you understand risks,

Patient initials _____

potential complications, limitations, and consequences of Botox injections. Additional information concerning Botox may be obtained from the package-insert sheets supplied by Allergan.

Bleeding and Bruising- It is possible, though unusual, to have a bleeding episode from a Botox injection. Bruising in soft tissues may occur. Serious bleeding around the eyeball during deeper Botox injections for crossed eyes (strabismus) has occurred. Should you develop post injection bleeding, it may require emergency treatment or surgery. Aspirin, anti-inflammatory medications, platelet inhibitors, anticoagulants, Vitamin E, ginkgo biloba, and other "herbs / homeopathic remedies" may contribute to a greater risk of a bleeding problem. Do not take these for ten days before or after Botox injections.

Damage to Deeper Structures- Deeper structures such as nerves, blood vessels, and the eyeball may be damaged during the course of injection. Injury to deeper structures may be temporary or permanent.

Corneal Exposure Problems - Some patients experience difficulties closing their eyelids after Botox injections and problems may occur in the cornea due to dryness. Should this rare complication occur, additional treatments, protective eye drops, contact lenses, or surgery may be necessary.

Dry Eye Problems- Individuals who normally have dry eyes may be advised to use special caution in considering Botox injections around the eyelid region.

Migration of Botox- A black-box warning is listed in the package insert of Botox cosmetic. This indicates that the effects of Botox may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. They have been reported to occur hours to weeks after injection. Swallowing and breathing difficulties can be life-threatening and there have been reports of death. These reports are more common in patients who have an underlying condition that would predispose them to these symptoms.

Drooping Eyelid (Ptosis) - Muscles that raise the eyelid may be affected by Botox, should this material migrate downward from other injection areas.

Double-Vision - Double-vision may be produced if the Botox material migrates into the region of muscles that control movements of the eyeball.

Eyelid Ectropion- Abnormal looseness of the lower eyelid can occur following Botox injection.

Other Eye Disorders - Functional and irritative disorders of eye structures may rarely occur following Botox injections.

Blindness- Blindness is extremely rare after Botox injections. However, it can be caused by internal bleeding around the eyeball or needle stick injury. In a period of 10 years of Botox administration, complications of blurred vision, retinal vein occlusion, and glaucoma have been reported in three patients. The occurrence of eye problems appears to be very rare.

Asymmetry- The human face and eyelid region is normally asymmetrical with respect to structural anatomy and function. There can be a variation from one side to the other in terms of the response to Botox injection.

Pain- Discomfort associated with Botox injections is usually short duration.

Allergic Reactions- As with all biologic products, allergic and systemic anaphylactic reactions may occur. Allergic reactions may require additional treatment.

Antibodies to Botox- Presence of antibodies to Botox may reduce the effectiveness of this material in subsequent injections. The health significance of antibodies to Botox is unknown.

Infection- Infection is extremely rare after Botox injection. Should an infection occur, additional treatment including antibiotics may be necessary.

Skin Disorders- Skin rash, itching, and swelling may rarely occur following Botox injection.

Neuromuscular Disorders- Patients with peripheral motor neuropathic disorders (amyotrophic lateral sclerosis, myasthenia gravis, motor neuropathies) may be at greater risk of clinically significant side effects from Botox.

Patient initials _____

Unsatisfactory Result - There is the possibility of a poor or inadequate response from Botox injection. Additional Botox injections may be necessary. Surgical procedures or treatments may be needed to improve skin wrinkles including those caused by muscle activity.

Long-Term Effects - Subsequent alterations in face and eyelid appearance may occur as the result of aging, weight loss, weight gain, sun exposure, pregnancy, menopause, or other circumstances not related to Botox injections. Botox injection does not arrest the aging process or produce permanent tightening of the eyelid region. Future surgery or other treatments may be necessary.

Pregnancy and Nursing Mothers- Animal reproduction studies have not been performed to determine if Botox could produce fetal harm. It is not known if Botox can be excreted in human milk. It is not recommended that pregnant women or nursing mothers receive Botox treatments.

Drug Interactions- The effect of Botox may be potentiated by aminoglycoside antibiotics or other drugs known to interfere with neuromuscular transmission.

Unknown Risks - The long-term effect of Botox on tissue is unknown. The risk and consequences of accidental intravascular injection of Botox is unknown and not predictable. There is the possibility of additional risk factors may be discovered.

HEALTH INSURANCE

Most health insurance companies exclude coverage for cosmetic surgical procedures and treatments or any complications that might occur from the same. Please carefully review your health insurance subscriber information pamphlet.

ADDITIONAL TREATMENT NECESSARY

There are many variable conditions in addition to risk and potential complications that may influence the long-term result of Botox injections. Even though risks and complications occur infrequently, the risks cited are the ones that are particularly associated with Botox injections. Other complications and risks can occur but are even more uncommon. Should complications occur, additional surgery or other treatments may be necessary. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained.

FINANCIAL RESPONSIBILITIES

It is unlikely that Botox injections to treat cosmetic problems would be covered by your health insurance. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome. Additional costs of medical treatment would be your responsibility should complications develop from Botox injections.

In signing the consent for this surgery/procedure, you acknowledge that you have been informed about its risk and consequences and accept responsibility for the clinical decisions that were made along with the financial costs of all future treatments.

Patient Name

Witness

Date

Patient initials_____